

## Antiretroviral medications\* to be stocked in hospital pharmacy

NOTE: These medications will be needed expediently for patient administration to reduce HIV mother to child HIV transmission (MTCT) in setting of positive maternal rapid HIV test in labor and delivery

Medication	How Supplied:	Storage Requirements:	Preparation:
<b>Zidovudine (Retrovir®) IV Infusion, 10 mg zidovudine/mL</b>  Manufacturer: GlaxoSmithKline	20-mL Single-Use Vial, Tray of 10	15-25 °C (59-77 °F) , protect from light.	Dilute prior to administration. After dilution, physically and chemically stable 24 hours at room temperature, 48 hours if refrigerated at 2-8 °C (26-46 °F).
<b>Zidovudine (Retrovir®) Tablets 300 mg</b>  Manufacturer: GlaxoSmithKline	Bottle of 60	15-25 °C (59-77 °F)	
<b>Zidovudine (Retrovir®) Syrup, 50 mg zidovudine/5mL</b> (clear colorless to pale yellow liquid)  Manufacturer: GlaxoSmithKline	240 mL bottle with child-resistant cap	15-25 °C (59-77 °F)	
<b>Lamivudine (Epivir®) 150mg tablets</b>  Manufacturer: GlaxoSmithKline	Bottle of 60 tablets, child-resistant closure	15-30 °C (59-86 °F)	
<b>Lamivudine (Epivir®) Oral solution, 10 mg lamivudine/1mL</b> (clear colorless to pale yellow liquid)  Manufacturer: GlaxoSmithKline	240 mL plastic bottle, child-resistant closure	25 °C (77 °F) USP Controlled Room Temperature	
<b>Combivir® (lamivudine 150mg/zidovudine 300mg) tablets</b>  Manufacturer: GlaxoSmithKline	Bottle of 60 tablets or unit dose pack of 120 tablets	2-30 °C (36-86 °F)	
<b>Nevirapine (Viramune®) Tablets 200 mg</b>  Manufacturer: Boehringer-Ingelheim	Bottle of 60 tablets	Storage requirements: 25 °C (77 °F) USP Controlled Room Temperature, excursions permitted to 15-30 °C (59-86 °F)	
<b>Nevirapine (Viramune®) Oral Suspension, 50 mg nevirapine/5mL</b> (off-white preserved suspension)  Manufacturer: Boehringer-Ingelheim	240 mL plastic bottle, child-resistant closure	25 °C (77 °F) USP Controlled Room Temperature, excursions permitted to 15-30 °C (59-86 °F)	

\*Medication information adapted from package inserts posted at [www.fda.gov](http://www.fda.gov) January 2007.