

Medical Record Number

Patient Name

Addressograph or Label

L13539

PROGRESS NOTE • POSITIVE RAPID HIV TEST
COUNSELING FORM

- Positive Rapid HIV result reported to patient
- Advised patient of the following:
 - Results are preliminary, and further testing is needed to confirm a true positive result.
 - There are interventions to decrease the risk of perinatal HIV transmission such as antiretroviral medications.
 - Risk of transmission without intervention is approximately 25% (1 in 4).
 - In the event that preliminary results are falsely positive (i.e. the rapid test is positive, but the confirmatory test returns as negative), current interventions would be unnecessary; the risk of HIV transmission with a positive preliminary result, however, outweighs the risks of these interventions.
 - Breastfeeding is not recommended until confirmatory testing is complete. Until then the patient may pump her breast milk to establish a milk supply.
- Recommendations
 - From rapid HIV back up physician: Name _____
 - From National Perinatal Hotline: 888-448-8765

Mode of delivery:

Pre-labor preliminary positive

- Recommend cesarean delivery.

Intrapartum preliminary positive

- Plan for anti-retrovirals and vaginal delivery with cesarean delivery for routine obstetric indications.

Antiretrovirals:

- IV AZT
- Other:

No AROM without attending approval.
No fetal scalp electrode or scalp sampling.
Pitocin to shorten labor if not progressing.

- Pediatric team notified
- Social Work notified

Name/Title _____ Pager: _____ Date _____ Time: _____