

Medical Record Number

Patient Name

Addressograph or Label

L13539

ORDERS • INTRAPARTUM PHYSICIAN ORDERS FOR PREGNANT PATIENTS WITH POSITIVE RAPID HIV TESTS

ALLERGIES _____

GENERAL Orders: (* = precautionary statement) (Mark "x" in box to order.)

Page Rapid HIV back up physician:

1. **Rahangdale (13182)**

2. **Daniels (13844)**

3. **Brodzinsky (13508)**

4. **MFM backup on call**

- Weigh patient.
- Labs: HIV viral load, CD4, LFTs, RPR, Type and screen, and CBC w/ diff.
- No fetal scalp electrode or scalp sampling
- No artificial rupture of membranes without approval of attending
- DO NOT USE Methergine for management of post-partum hemorrhage/uterine atony unless alternative treatments are not available (*potential for excessive vasoconstriction if given with protease inhibitor or with non-nucleoside reverse transcriptase inhibitors)
- Administer pitocin per protocol to shorten duration of labor, especially if membranes are ruptured.
- Notify Newborn nursery team and Pediatric ID fellow on call.
- Alert perinatal social worker.

ANTIRETROVIRAL (ARV) Orders for labor: (* = precautionary statement)

- Zidovudine (AZT) Loading Dose = _____ mg (2 mg/kg) IV over one (1) hour. Follow by continuous infusion of _____ mg/hr (1 mg/kg/hr) until umbilical cord is clamped.**

Pharmacy to prepare continuous infusion of Zidovudine 1000 mg in D5W 250 ml to run at _____ ml/hr = 1 mg/kg/hr.

Body Weight = _____ kg

***DO NOT RUN AZT IN SAME IV LINE AS MAGNESIUM SULFATE**

- Lamivudine 150 mg PO, 1 tab BID
- Nevirapine 200 mg PO, 1 tab once daily
- Other _____

OTHER MEDICATIONS:

- Other _____

POSTPARTUM

- Discontinue IV AZT when cord clamped
- Page Rapid HIV back up physician for postpartum antiretroviral orders

Date: _____ Time: _____ Provider: _____ / _____ Pager # _____
Print name Signature Title

Date: _____ Time: _____ RN signature: _____